

DSSY implementation on deal. II

Imbunm Kim

Seoul National University *ibkim11@gmail.com*

DSSY Nonconforming Quadrilateral Finite Element
(RAIRO Anam. Numer. 33, 1999, Douglas-Santos-Sheen-Ye)

On a reference domain $\hat{Q} := [-1, 1]^2$, define

$$DSSY(\hat{Q}) = \text{span}\{1, \hat{x}, \hat{y}, \theta_l(\hat{x}) - \theta_l(\hat{y})\}, \quad l = 1, 2$$

$$\text{where } \theta_l(x) := \begin{cases} x^2 - \frac{5}{3}x^4, & l = 1 \\ x^2 - \frac{25}{6}x^4 + \frac{7}{2}x^6, & l = 2 \end{cases} \quad (1)$$

The reference four DSSY degrees of freedom are given at the midpoints.

Let $F_Q : \hat{Q} \rightarrow Q$ be a bijective affine transformation. Then

$DSSY(Q) = \{\hat{v} \circ F_Q^{-1} \mid \hat{v} \in DSSY(\hat{Q})\}$. Then DSSY FEM is defined by

$DSSY^h = \{v \in L^2(\Omega) \mid v|_Q \in DSSY(Q) \text{ for all } Q \in$

$\tau_h; v \text{ is continuous at the midpoint of each interior edge}\}$

DSSY implementation on deal.II




Figure: (step3) Q_1 CFE / DSSY NCFE solution (step4) Q_1 CFE / DSSY NCFE solution

Thing to do : Apply DSSY to other test problems ; Changes should be made in nonconforming sense