

Jaeryun YIM from Seoul National University

My works:

- 1  $P_1$  Nonconforming FE
  - piecewise linear function on quadrilateral element
  - continuous at midpoint of each edge
- 2 Residual Based A Posteriori Error Estimator for  $P_1$  NCFE

$$\eta^2 := \sum_K h_K^2 \underbrace{\|f + \operatorname{div} \nabla u_h^{nc}\|_{0,K}^2}_{\text{cell residual}} + \sum_E h_E \left( \underbrace{\|J_{E,\nu}\|_{0,E}^2}_{\text{normal jump}} + \underbrace{\|J_{E,\tau}\|_{0,E}^2}_{\text{tangential jump}} \right) \quad (1)$$


$$\text{where } J_{E,\nu} := \begin{cases} \nu_1 \cdot \nabla u_h^{nc}|_{K_1} + \nu_2 \cdot \nabla u_h^{nc}|_{K_2} & E \in \mathcal{E}(\Omega) \\ g - \nu \cdot \nabla u_h^{nc} & E \in \mathcal{E}(\Gamma_N) \\ 0 & E \in \mathcal{E}(\Gamma_D) \end{cases} \quad (2)$$

$$J_{E,\tau} := \begin{cases} \tau_1 \cdot \nabla u_h^{nc}|_{K_1} + \tau_2 \cdot \nabla u_h^{nc}|_{K_2} & E \in \mathcal{E}(\Omega) \\ 0 & E \in \mathcal{E}(\Gamma_N) \\ \tau \cdot (\nabla u_D - \nabla u_h^{nc}) & E \in \mathcal{E}(\Gamma_D) \end{cases} \quad (3)$$

Consider  $-\Delta u = 0$  in  $\Omega = [-1, 1]^2 \setminus [0, 1]^2$

Exact solution in polar coordinate

$$p(r, \theta) = r^\alpha \sin(\alpha(\theta - \pi/2)) \in H^{1+\alpha}(\Omega), \quad \alpha = 2/3 \quad (4)$$



**Figure :** (Tops)  $P_1$  NCFE Solution w/ tangential term: (L) After 1 refinement, (R) After 7  
 (Bottoms) Grid after 7 refinements:  
 (L)  $Q_1$  CFE, (C)  $P_1$  NCFE w/o tangential term, (R)  $P_1$  NCFE w/ tangential term

Further works:

- tangential jump term on Dirichlet boundary edges
- coefficient consideration for the general elliptic problem
- a posterior error estimator for NCFE to another problems