Handling non-matching methods between independent distributed grids: step-70

Luca Heltai¹, Bruno Blais²

¹International School for Advanced Studies ²Polytechnique Montréal





26 May 2020 - deal.II Online Workshops

Outline



Introduction

- Motivations
- Everything reduces to point evaluation
- We need a way to track points: particles

Fluid structure interaction

- Matching solids points into fluid cells
- Insert global particles
- Keeping track of ownership (LAC)

Putting it all together: Step-70

- Nitsche method
- A classical experiment
- Results: impellers in 2 and 3 dimensions

Plan



Introduction

- Motivations
- Everything reduces to point evaluation
- We need a way to track points: particles

- Matching solids points into fluid cells
- Insert global particles
- Keeping track of ownership (LAC)

- Nitsche method
- A classical experiment
- Results: impellers in 2 and 3 dimensions

Mixing

- Critical industrial application
- Often a severe bottleneck that leads to waste
- Costly and difficult to investigate experimentally
- × Not trivial so simulate



Simulation of mixing

Rotation of the geometry prevents the use of ALE techniques. Requires the use of specialized techniques:

- **X** Change of reference frame: Severe limitations on the geometry
- ✗ Sliding mesh: Requires coupling the boundaries of two grids
- ✓ Non-matching methods: immersed boundaries, fictious domains, etc.

Non-matching methods can also be used for a much broader range of applications:

- ✓ Fluid-Structure interaction
- ✓ Overlaping geometries (e.g. Planetary mixers)



Graphical summary of non-matching methods



- Different properties on different domains
- ✓ Independent discretizations (FEM, BEM, FV, FD, etc.)
- Different physics on different domains (i.e., fluids and solids)
- Possibly different intrinsic scales
- We need a numerically efficient way to transfer information between the domains

We need a way to track points: particles

Particles are an ideal tool to keep track of the motion of our embedded domain:

- ✓ They can readily move from processor to processor and can be exchanged using some-to-some communications.
- ✓ They already contain a very efficient mechanism to detect in which cell they lie and what is their reference location.
- ✓ An arbitrary amount of properties can be attached to them.



1 2 GridGenerator::generate_from_name_and_arguments(
 fluid_tria, par.name_of_fluid_grid,
 par.arguments_for_fluid_grid);













Plan



- Motivations
- Everything reduces to point evaluation
- We need a way to track points: particles

Fluid structure interaction

- Matching solids points into fluid cells
- Insert global particles
- Keeping track of ownership (LAC)

Putting it all together: Step-70

- Nitsche method
- A classical experiment
- Results: impellers in 2 and 3 dimensions

Insert global particles in distributed environments

Generating the particles of the embedded domain is quite challenging in parallel distributed environments:

- The two domains do not match
- The points of the embedded domain can lie in a cell which belongs to another processor



Requires a global insertion mechanism to create particles from an :

```
std::vector<Points<dim>>
```

where each processor owns such a (possibly non-empty) vector.

1

Insert global particles

1

2

3

4

1

1

1

✓ The vector containing the location of the particles created by the processor. These particle do not need be located in a cell in that processor.

```
std::vector<Points<dim> > points_vector
```

✓ An (n_{proc}) size vector containing a vector of bounding boxes

std::vector<std::vector<BoundingBox<spacedim>>
global_bounding boxes

A vector of vectors of particle properties

std::vector<std::vector<double> >
 particle_properties

boost::rtree of Bounding boxes surrounding cells

A boost::rtree of bounding boxes allows the identification of the cells in which the particles lie in $\mathcal{O}(n_{particles} \log(m_{boxes}))$ time.



Luca Heltai¹, Bruno Blais²

Step 70

boost::rtree of Bounding boxes in parallel



Bounding boxes

1

2

Generating the bounding boxes is relatively easy.

```
1
  std::vector<BoundingBox<spacedim>> all_boxes;
2
  all_boxes.reserve(fluid_tria.n_locally_owned_active_cells());
3
  for (const auto cell : fluid_tria.active_cell_iterators())
     if (cell->is_locally_owned())
4
5
       all_boxes.emplace_back(cell->bounding_box());
6
7
  const auto tree = pack_rtree(all_boxes);
8
  const auto local_boxes =
9
     extract_rtree_level(tree,
        par.fluid_rtree_extraction_level);
0.
.1
  global_fluid_bounding_boxes =
2
    Utilities::MPI::all_gather(mpi_communicator,
        local_boxes);
```

and bounding boxes can now even be written and visualized!

 BoundingBoxDataOut<dim> data_out; data_out.build_patches(my_bounding_box);

 Luca Heltai¹, Bruno Blais²

Step 70
26 May 2020 - deal.II (Zoom)
17/31

Insert global particles

Finally, the key steps in the global particle insertion:

- Gather the number of particles to be added for each processors (all_gather)
- Identify on which cell and on which processor the particle are located (distributed_compute_point_location)
- Gather the properties of the particles from the original owner of the points
- Generate the particles with their properties and unique id

Outcome: Flexible particle insertion mechanisms

- Insertion at DOF support points
- Insertion at the quadrature points of a triangulation



As long as you have a triangulation, you can now use it to insert particles.

19/31

Keeping track of ownership: Linear algebra coupling



- We keep around two IndexSet objects, to identify locally owned particles coordinates (blue dots) ...
- ... and locally relevant particle coordinates (brown dots), i.e., coordinates of particles that fall within the current fluid domain, owned by someone else

Luca Heltai¹, Bruno Blais²

Initializing ownership

1

2

3

1 2 3

Either based on where particles end up in the first step:

```
locally_owned_tracer_particle_coordinates =
    tracer_particle_handler.locally_relevant_ids().tensor_produc
    complete_index_set(spacedim));
```

... or based on solid ownership (as in the previous image). Then at each step, keep track of locally relevant ones in the same way:

<pre>locally_relevant_tracer_particle_coordinates =</pre>	
<pre>tracer_particle_handler.locally_relevant_ids().tensor_pro</pre>	du
<pre>complete_index_set(spacedim));</pre>	

And initialize a parallel MPI vector with it...

	relevant_tracer_particle_displacements.reinit(
	<pre>locally_owned_tracer_particle_coordinates,</pre>
ĺ	<pre>locally_relevant_tracer_particle_coordinates,</pre>
ĺ	<pre>mpi_communicator);</pre>

... that can be used as a base for the solid equations.

Updating velocities

1

1 2 Interpolating from fields to particles:

1	Particles::Utilities::interpolate_field_on_particles(I
2	fluid_dh,	I
3	<pre>tracer_particle_handler,</pre>	I
4	locally_relevant_solution,	I
5	<pre>tracer_particle_velocities,</pre>	I
6	fluid_fe->component_mask(FEValuesExtractors::Ve	ct
7		I
8	<pre>tracer_particle_velocities *= time_step;</pre>	I

... update the relevant IndexSet, and copy locally (read only!) the velocities of all the particles that fall within the locally owned fluid domain (MPI communication happens in the background):

```
relevant_tracer_particle_displacements =
   tracer_particle_velocities;
```

... and transfer information back to the particles:

```
tracer_particle_handler.set_particle_positions(
            relevant_tracer_particle_displacements);
```

Plan



- Motivations
- Everything reduces to point evaluation
- We need a way to track points: particles

Fluid structure interaction

- Matching solids points into fluid cells
- Insert global particles
- Keeping track of ownership (LAC)

3 Putting it all together: Step-70

- Nitsche method
- A classical experiment
- Results: impellers in 2 and 3 dimensions

Physics solved

In step-70, we solve the Stokes equations for a creeping flow (i.e. a flow where Re \rightarrow 0) and a no-slip boundary condition is applied on a moving *embedded domain* Γ associated with an impeller.

This leads to the following differential problem: given a sufficiently regular function g on Γ , find the solution (\mathbf{u}, p) to

$$-\Delta \mathbf{u} + \nabla p = 0, \tag{1}$$

$$-\nabla \cdot \mathbf{u} = \mathbf{0},\tag{2}$$

$$\mathbf{u} = \mathbf{g} \text{ in } \Gamma, \tag{3}$$

$$\mathbf{u} = 0 \text{ on } \partial \Omega.$$
 (4)

Imposing the motion of the impeller

Two scenarios occur when enforcing conditions on the embedded domain:

- The geometrical dimension 'dim' of the embedded domain Γ is the same of the domain Ω ('spacedim'), In this case, the imposition of the Dirichlet boundary boundary condition on Γ is done through a volumetric penalization.
- The embedded domain Γ has an intrinsic dimension 'dim' which is smaller than that of Ω ('spacedim'), thus its spacedim-dimensional measure is zero; for example it is a curve embedded in a two dimensional domain, or a surface embedded in a three-dimensional domain. In this case, the boundary condition is imposed weakly on Γ by applying the Nitsche method.

For both cases, a possible resulting formulation is:

$$(\nabla \mathbf{v}, \nabla \mathbf{u})_{\Omega} - (\operatorname{div} \mathbf{v}, \mathbf{p})_{\Omega} - (\mathbf{q}, \operatorname{div} \mathbf{u})_{\Omega} + \beta(\mathbf{v}, \mathbf{u})_{\Gamma} = \beta(\mathbf{v}, \mathbf{g})_{\Gamma}.$$

< 同 ト < 三 ト < 三 ト

Assembly of the Nitsche terms – Part I

Looping over all particles, cell-wise, and getting information from particles:

```
auto particle = solid_particle_handler.begin();
while (particle != solid_particle_handler.end())
  Ł
    const auto &cell =
       particle->get_surrounding_cell(fluid_tria);
    const auto pic =
       solid_particle_handler.particles_in_cell(cell);
    Assert(pic.begin() == particle,
       ExcInternalError());
    for (const auto &p : pic)
      Ł
        const auto &ref_q
           p.get_reference_location();
        const auto &real_q = p.get_location();
        const auto &JxW = p.get_properties()[0];
        . . .
```

1

2

3

4

5

6

7

8

9

.0

.1

2

A B A B A B A B A B A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A
 B
 A

Assembly of the Nitsche terms - Part II

Assemble the penalization matrix:

1

2

3

4

5

6 7

8

9

10

11

12

13

14

15

16

17

18

19

```
for (unsigned int i = 0; i < dofs_per_cell; ++i) {</pre>
    const auto comp_i =
      fluid_fe->system_to_component_index(i).first;
    if (comp_i < spacedim) {</pre>
        for (unsigned int j = 0; j < dofs_per_cell; ++j)</pre>
            const auto comp_j =
              fluid_fe->system_to_component_index(j).first;
            if (comp_i == comp_j)
              local_matrix(i, j) +=
                 penalty_parameter * par.penalty_term *
                 fluid_fe->shape_value(i, ref_q) *
                 fluid_fe->shape_value(j, ref_q) * JxW;
          3
        local_rhs(i) += penalty_parameter * par.penalty_term *
                         solid_velocity.value(real_q, comp_i) *
                         fluid_fe->shape_value(i, ref_q) * JxW;
      }
  }
```

- 34

イロト イポト イヨト イヨト

The problems: Mixing in the creeping flow regime

- ✓ Famous experiment by Sir G.I. Taylor using coloured dyes
- ✓ Multiple similar experiments on Youtube



2D results



3.0
2.5
2.0
2.0
1.5
M dobuit of the second sec

29 / 31



Impeller Tracer Particles

Luca Heltai¹, Bruno Blais²

Step 70

26 May 2020 - deal.II (Zoom)

3D results



æ

< □ > < □ > < □ > < □ > < □ >

Conclusions

Step-70 demonstrates a large variety of concepts

- Global insertion of particles
- ✓ Linear algebra coupling between particles and DOFS
- Non-matching methods
- ✓ Post-processing of particles using the new Particles::DataOut

It opens up a large range of possibilities for the solution of:

- Massively parallel immersed boundary problems
- Fluid structure interaction
- Chimera grids
- ✓ and many others...

It also was a very fun collaborative work!